


# ArborTags



The leaves are finely sawtoothed, thickened, and have often a blunt end. They are shiny green above with prominently sunken veins. The Rusty Blackhaw has rust colored hairs on the underside.

[www.arborworks.org](http://www.arborworks.org)

*Viburnum rufidulum*

**Rusty Blackhaw**

**Rusty Nannyberry**

*Viburnum rufidulum*

**Rusty Blackhaw**

**Rusty Nannyberry**



The flowers are themselves tiny, but cluster in a 3-5 inch aggregate. When they bloom in the spring their showy white display is visible from afar. Rust colored hairs are visible amidst the white blossoms.

[www.arborworks.org](http://www.arborworks.org)

©1996 American Association of Amateur Arborists



The fruit is a small blue-black berry with a whitish frosting. A favorite of songbirds the berries are slightly sweet

[www.arborworks.org](http://www.arborworks.org)

*Viburnum rufidulum*

**Rusty Blackhaw**

**Rusty Nannyberry**

*Viburnum rufidulum*

**Rusty Blackhaw**

**Rusty Nannyberry**

The Blackhaw is often a bush or a small tree with a short trunk. The rough gray bark breaks up into small plates and appears similar to the bark of a flowering dogwood. Also like the dogwood the leaves form opposite one another.

[www.arborworks.org](http://www.arborworks.org)